

Dairy Fat Consumption and Mortality due to Cardiovascular Disease. Does Source of Fats Matter?

Paula Moliterno^{1,*}, Florencia Cerruti²

¹Escuela de Nutrición, Universidad de la República. Montevideo, Uruguay

²Universidad Católica del Uruguay. Montevideo, Uruguay

Abstract This study analyzed data for correlation between dairy fat availability for consumption and cardiovascular mortality, in adults of both sexes from 35 to 74 years of age, across a selection of countries. In addition we focused on the dietary levels of vegetable and animal fats in countries with high and low cardiovascular mortality. Mortality statistics were taken from the WHO, 2014. Data pertaining to fat availability for consumption was obtained from Food Balance Sheets (FAOStat, 2011). Consumption patterns of animal and vegetal fats were analyzed for countries with high and low cardiovascular mortality. Evolution of dairy fat consumption has changed over time. Cardiovascular mortality showed high rates in countries within the region. Countries with low cardiovascular mortality showed higher levels of vegetable fat consumption, while countries with high mortality rates showed higher levels of animal fat consumption. No particular pattern was observed for dairy fat. This study found no significant correlation between dairy fat consumption and cardiovascular mortality.

Keywords Dairy Fat, Cardiovascular Disease, Mortality, Vegetable and Animal Fats

1. Introduction

Cardiovascular disease (CVD) is responsible for approximately 17 million annual deaths greatly affecting countries of low and medium income, determined in 2008 the main cause of mortality in adults younger than 60 years old [1]. The World Health Organization (WHO) estimates that the number of deaths caused by this disease will grow 23.3 million by 2030, continuing as the leading cause of death globally [2].

Unhealthy diet has been defined by the WHO as a modifiable risk factor and fundamental determining in the incidence of CVD [3], reason why the lipid fraction of the diet has traditionally been involved in its management and risk prevention [4]. International statistics indicate a strong correlation between the consumption of saturated fat (from animal products) and mortality caused by coronary heart disease [5].

Evidence shows that high intake of saturated fat (SF) and industrial trans fatty acids is associated to a greater incidence of these diseases [6], so the recommendation to diminish its consumption has become a priority health measure to reduce the risk of CVD [7-9].

Traditionally, dairy products have played a fundamental

role in human nutrition associated to their composition rich in nutrients; however, dairy fat has a varied fatty acid composition, highlighting the saturated palmitic fatty acid with proven metabolic effect on cardiovascular health. An ecologic study on year 1979 showed a strong correlation between the availability of dairy fat and mortality by cardiovascular disease ($r= 0,75$) [10].

These results along with those gathered by the Seven Countries Study [11] have determined a recommendation on the inclusion of low-fat dairy in promoting cardiovascular health. Nonetheless, lately, evidence available shows contradicting and inconclusive results related to the consumption of dairy products and risk and mortality due to CVD [12-15].

In this context, we proposed the hypothesis that a higher availability of dairy fat for human consumption is correlated to higher rates of cardiovascular mortality, since dairy products constitute food which saturated fat has fatty acids known for its implication in the development of CVD.

The main objective of this study was to observe the evolution in the consumption of dairy fat and the availability of vegetable and animal fat in a group of countries where the rates of cardiovascular mortality cover a broad spectrum. Intended to provide new information relevant to elucidate the contradictions described in previous paragraphs, the relation between availability of dairy fat for consumption and cardiovascular mortality rates in countries with extreme cardiovascular mortality was analyzed.

* Corresponding author:

pmoliterno@nutricion.edu.uy (Paula Moliterno)

Published online at <http://journal.sapub.org/fph>

Copyright © 2016 Scientific & Academic Publishing. All Rights Reserved

2. Material and Method

A descriptive, observational and cross-sectional study was designed.

Data of availability of dairy fat, animal fat and vegetable fat for human consumption was obtained through Food Balance Sheets of FAO (FAOStat, 2011) for the period of 1970 to 2011 [16]. This instrument allows determining the apparent national consumption of each food, presenting the structure of the main food supply of a country during a reference period. Based on this information, animal fat are subcategorized in meat fats and dairy fats. Meat fat constituted the animal fat except dairy fat.

Data of 24 countries of different continents were added considering those included in the Turpeinen study of 1979; United States and Canada, Italy, Portugal, France, England, Netherlands, Switzerland, Spain, Poland, Finland, Romania, Sweden, Austria, Norway and Germany; Japan; Australia and New Zealand. In order to complement the analysis with Latin American countries that were not included in the study mentioned, data from Uruguay, Argentina, Chile, Brazil and Costa Rica were included.

For cardiovascular mortality analysis, data published by the WHO in 2014 [17] was used. Age-standardized cardiovascular mortality rates were analyzed for adult age (35 to 74 years old). Additionally, specific-sex mortality rates were analyzed. Cardiovascular mortality causes were considered as all below the item “circulatory system diseases” in the International Classification of Diseases (ICD-10) [18].

For the study of the consumption of animal and vegetal fats, 10 countries in the extreme spectrum of mortality rates were selected. Based on prior, low mortality rates were considered fewer than 100 (each 100.000) and high mortality, rates higher or equal to 100.1 (each 100.000).

To study the association between the availability of dairy fat and mortality due to CVD in 24 chosen countries, the statistical Pearson correlation coefficient test was applied.

All analyses were conducted with the use of SPSS statistical software.

3. Results

Mortality rates due to CVD are shown comparatively in Figure 1. The difference in cardiovascular mortality was more than 5 times between the first place Rumania (436/100.000) and the last one France (80.4/100.000). Statistics indicate that Rumania (436 each 100.000) and Poland (294.2 each 100.000) are among the countries which cardiovascular mortality rates are higher. In Latin America, Brazil (249 each 100.000), Argentina (198.5 each 100.000) and Uruguay (176.9 each 100.000) were placed with higher cardiovascular mortality rates.

Dairy fat availability in America has varied from 15.6

grams (g) per capita (1970-1975) to 17.4 g per capita in 2011. In Asia, it is under 10.0 g of dairy fat per capita, varying from 3.2 g per capita (1970-1975) to 7.9 g in 2011. Levels of dairy fat availability in Europe varied from 29.5 g per capita (1970-1975) to 23.4 g per capita in 2011.

In the countries with high cardiovascular mortality, availability of vegetal, meat and dairy fats for consumption, considered individually, was placed under 60.0 g, whilst in the countries with a low cardiovascular mortality this was placed under 85.0 g. This is illustrated in Figure 2.

Table 1 shows the variations in the availability of dairy fat in the year 1973 and its correspondent in year 2009 for the selection of countries included in [10]. Likewise, cardiovascular mortality rates in adult men in the countries previously mentioned are shown comparatively for year 1973 and 2009 on Figure 3.

Table 1. Variations on Availability of Dairy Fat (g) for Human Consumption between 1973 and 2009 in a Countries Selection. FAOStat

Country	1973	2009	Variation (%)
Japan	5,2	7,0	34,6
Portugal	8,8	18,4	109,1
Spain	14,0	18,8	34,3
Italy	20,2	25,6	26,7
USA	23,7	26,8	13,1
Austria	31,6	30,1	-4,7
Germany	32,7	31,8	-2,8
Canada	32,8	24,0	-26,8
Sweden	35,1	36,2	3,1
Australia	35,9	26,8	-25,3
England	38,1	28,1	-26,2
Poland	41,1	26,8	-34,8
Norway	41,1	29,7	-27,7
Switzerland	41,2	38,4	-6,8
Finland	60,0	39,3	-34,5

On the countries where male cardiovascular mortality increased, variations in the availability of dairy fat consumption were observed. Japan increased cardiovascular mortality on 194%, while the availability of dairy fat in the same period grew on 35%. In Portugal, cardiovascular mortality rose 30% along with the rise of 109% in the availability of dairy fat. Poland increased mortality due to CVD by 187% with a decline of 35% in the availability of dairy fat. Other countries showed a decline in cardiovascular mortality between 1973 and 2009.

From the analysis of the association between the availability of dairy fat and CVD mortality it was found that there is no statistically significant correlation in men ($r = 0.004$ $p = 0.77$) (Figure 4) nor in women ($r=0.03$ $p=0.42$).




Figure 1. Cardiovascular mortality among adults (35-74 years old), both gender, in a selection of 24 countries. WHO 2009




Figure 2. Availability for human consumption of major food fats in 10 countries with high and low cardiovascular mortality, respectively. V = vegetable fats, M = meat fats, D = dairy fats




Figure 3. Comparison of mortality due to CVD among adult men (35-37 years old) for a countries selection. WHO 1973, 2009

4. Discussion

The main find of this study was that currently, no statistically significant correlation exists between availability of dairy fat for consumption and mortality due to CVD, unlike past findings ($r = 0,75$ $p < 0,001$) [10], which may not allow to confirm the hypothesis at first formulated. At the present time, an increase in availability of dairy fat for consumption in adult men would not be statistically associated with increases in mortality rates due to CVD.

Moreover, changes in location of countries at the correlation line were observed, showing differences among those two variables over time. In 1979, countries such as Finland and New Zealand of important dairy production were located at the upper extreme of the correlation line, whilst Asian countries like Japan and the Mediterranean countries were located on the lower extreme.

Currently, Finland is located among countries with major availability of dairy fat for consumption and lower mortality rate. Such variations could be related to successful community-based intervention actions that have been carried out since 1971 in order to reduce the problem of CVD in Nor Karelia, extending nationwide, with emphasis on unhealthy diet as one of the fundamental risk factor [19].

On the other hand, Japan showed lower cardiovascular mortality rates and modest availability of dairy fat for consumption, being vegetable fats the predominant type of food fats. However, the analysis of variation of male mortality rates over time showed an important increase which occurs with a non proportional increase of availability of dairy fat, coinciding with increasing evolution of availability of dairy fat in Asian region.

Uruguay, cattle country, valued for the quality of its dairy

products, showed availability of dairy fat among the modest and leading mortality rates due to CVD. The results of this study show that evolution of cardiovascular mortality over time does not correlate proportionally with evolution of availability of dairy fat for consumption.

In this regard, it would not be appropriate to draw conclusions based on a single risk factor such as dairy fat for its rich SF content, coinciding with results published by O'Sullivan et al. that when examining consumption of dairy products as a natural and key food group typically high in this fatty acids found no significant statistical association with mortality due to CVD [RR 0.87 (0.62; 1.20)] [20]. Likewise, Soedamah-Muthu et al. found no significant association between milk consumption and total mortality [RR per 200 ml/d 0.99 (0.95; 1.03)] [21].

The analysis of consumption patterns of meat, vegetable and dairy fats in countries with high and low cardiovascular mortality did not allow categorical statements such as ones observed in the past [10]. Even though countries with low mortality showed higher availability of vegetable fat and countries with high mortality exhibit higher availability of animal fat for consumption, no particular pattern was observed for dairy fat.

It is essential to consider dairy fat consumption in the global context of the evolution of different CVD risk factors. It is a complex nutrient whose beneficial effects have been proven on cardiovascular health. A meta-analysis conducted by Elwood et al., showed a reduction of 10% to 15% in the incidence of coronary heart disease in subjects who reported greater consumption of milk compared to those of lower consumption [14].

Another meta-analysis subsequently published by the same group showed decreased risk of coronary heart disease

5. Conclusions

In this study we found no statistically significant correlation between availability of dairy fat for consumption and mortality due to CVD. The evolution of cardiovascular mortality over time and its multi factorial nature makes necessary comprehensive strategies for management of cardiovascular health. Dairy foods are complex and many mechanisms have been described by which their consumption may manage cardiovascular health, regardless of fat content.

ACKNOWLEDGEMENTS

We would like to thank José Boggia MD, PhD for his suggestions and review of the manuscript. We acknowledge publication support by the School of Nutrition, University of the Republic, Uruguay.

REFERENCES

- [1] World Health Organization. Global status report on noncommunicable diseases. Geneva 2011.
- [2] Mathers C, Loncar D. Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med.* 2006; 3(11).
- [3] Pan American Health Organization. Non-communicable diseases in the Americas: Lets built a healthier future. Washington D.C 2011. (In spanish).
- [4] Srinath R, Katan M. Diet, nutrition and the prevention of hypertension and cardiovascular diseases. *Public Health Nutr.* 2004; 7: p. 167-186.
- [5] Siri-Tarino PW, Sun Q, Hu FB, Krauss RM. Saturated fat, carbohydrate, and cardiovascular disease. *Am J Clin Nutr.* 2010; 91: p. 502-509.
- [6] Michas G, Micha R, Zampelas A. Dietary fats and cardiovascular disease: Putting together the pieces of a complicated puzzle. *Atherosclerosis.* 2014; 234: p. 320-328.
- [7] Food and Agriculture Organization of the United Nations. Fats and fatty acids in human nutrition: Report of an expert consultation. FAO Food and Nutrition paper N° 91. Geneva. November; 2008.
- [8] World Health Organization. Diet, nutrition and prevention of chronic diseases. WHO Technical Report Series No 916. Geneva. 2003.
- [9] Eckel RH, Jakicic JM, Ard JD, et al. 2013 AHA/ACC Guideline on Lifestyle Management to Reduce Cardiovascular Risk. A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation.* 2013.
- [10] Turpeinen O. Effect of cholesterol-lowering diet on mortality from coronary heart disease and other causes. *Circulation.* 1979; 59: p. 1-7.
- [11] Verschuren W, Jacobs D, Bloemberg B, et al. Serum Total Cholesterol and Long-term Coronary Heart Disease Mortality in Different Cultures. Twenty-five-year Follow-up of the Seven Countries Study. *JAMA.* 1995 July; 274; 2: p. 131-136.
- [12] Huth P, Park K. Influence of dairy product and milk fat consumption on cardiovascular disease risk: a review of the evidence. *Adv Nutr.* 2012; 3: p. 266-285.
- [13] Rice BH. Dairy and cardiovascular disease: a review of recent observational research. *Curr Nutr Rep.* 2014; 3: p. 130-138.
- [14] Elwood PC, Givens DI, Beswick A, Fehily A, Pickering JE, Gallacher JE. The survival advantage of milk and dairy consumption: an overview of evidence from cohort studies of vascular diseases, diabetes and cancer. *J Am Coll Nutr.* 2008; 27: p. 723S-734S.
- [15] Elwood PC, Pickering JE, Givens DI, Gallacher JE. The consumption of milk and dairy foods and the incidence of vascular disease and diabetes: an overview of the evidence. *Lipids.* 2010 April; 45: p. 925-939.
- [16] Food and Agriculture Organization of the United Nations. FAOSTAT. Available from: <http://faostat3.fao.org> (1 July 2014). Rome, 2014.
- [17] World Health Organization. WHO Mortality Database. Available from: http://www.who.int/healthinfo/statistics/mortality_rawdata/en/index.html (7 July 2014) Department of Health Statistics and Information System. Geneva, Switzerland; 2013.
- [18] World Health Organization. ICD-10. International Classification of Diseases and related health problems. 10th Revision; 1995.
- [19] Puska P. Main Outline of The North Karelia Project. In Puska P, Vartiainen E, Laatikainen T, Jousilahti P, Paavola M. The North Karelia Project: From North Karelia to National Action. Helsinki: Helsinki University Printing House; 2009. p. 19-25.
- [20] O'Sullivan T, Hafekost K, Mitrou F, Lawrence D. Food sources of saturated fat and the association with mortality: A meta-analysis. *Am J Public Health.* 2013 September; 103: p. e31-e42.
- [21] Soedamah-Muthu SS, Ding EL, Al-Delaimy WK, et al. Milk and dairy consumption and incidence of cardiovascular diseases and all-cause mortality: dose-response meta-analysis of prospective cohort studies. *Am J Clin Nutr.* 2011; 93: p. 158-171.
- [22] Patterson E, Larsson SC, Wolk A, Akesson A. Association between dairy food consumption and risk of myocardial infarction in women differs by type of dairy food. *J Nutr.* 2013; 143: p. 74-79.
- [23] Nestel PJ, Chronopoulos A, Cehun M. Dairy fat in cheese raises LDL cholesterol less than that in butter in mildly hypercholesterolaemic subjects. *Eur J Clin Nutr.* 2005; 59: p. 1059-1063.
- [24] Alonso A, Nettleton J, Ix J, et al. Dietary Phosphorus, Blood Pressure, and incidence of hypertension in the Atherosclerosis Risk in Communities Study and the Multi-Ethnic Study of Atherosclerosis. *Hypertension.* 2010; 55: p. 776-784.
- [25] Aleixandre A, Miguel M, Muguera B. Antihypertension peptides derived from milk and egg proteins. *Nutr Hosp.* 2008; 23: p. 313-318. (In spanish).